



Vaccination Schedule for Broiler Chickens

Pin this on your poultry house wall or keep it with your farm records. Check off each vaccine as it's given, and fill in the actual date.

	DISEASE TARGET	VACCINE TYPE	ROUTE	GIVEN	DATE GIVEN
Day 1 (Hatchery)	Marek's Disease	HVT / Rispens strain	Subcutaneous injection	☐	_____
Day 1	Newcastle Disease (ND)	Live vaccine (B1 / La Sota)	Coarse spray or eye drop	☐	_____
Day 1 or 5-7	Infectious Bronchitis (IB)	Live vaccine (Massachusetts)	Coarse spray or eye drop	☐	_____
Day 10-14	Gumboro / IBD	Intermediate strain	Drinking water	☐	_____
Day 18-21	Newcastle Disease (Booster)	La Sota strain	Drinking water	☐	_____
Day 18-21	Infectious Bronchitis (Booster)*	Live vaccine	Drinking water or spray	☐	_____
Day 24-28	Gumboro (Booster)*	Intermediate / plus strain	Drinking water	☐	_____

***Optional / high-risk zones only.** Exact timing can shift based on regional disease pressure and hatchery practice. Always confirm your schedule with a local poultry veterinarian before applying it to your flock.

VACCINE STORAGE REMINDERS

- Store at 2°C–8°C (refrigerator)
- Do NOT freeze
- Protect from sunlight
- Use immediately after mixing
- Transport in a cool box with ice packs

ADMINISTRATION QUICK TIPS

- Thirst birds 1-2 hrs before water vaccines
- Use cool, dechlorinated water only
- Add skim milk powder to protect live virus
- Never vaccinate sick or stressed birds
- Keep full biosecurity on vaccination day

FARM NOTES